




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 209/42, 215/38, 215/40, 209/08, A61K 31/40, 31/47, 31/55		A1	(11) International Publication Number: WO 94/15919
			(43) International Publication Date: 21 July 1994 (21.07.94)
(21) International Application Number: PCT/GB93/02660		(74) Agents: BROWN, Keith, John, Symons et al.; Wyeth Laboratories, Huntercombe Lane South, Taplow, Maidenhead, Berkshire SL6 0PH (GB).	
(22) International Filing Date: 24 December 1993 (24.12.93)			
(30) Priority Data: 9300195.6 6 January 1993 (06.01.93) GB		(81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): JOHN WYETH & BROTHER LIMITED [GB/GB]; Huntercombe Lane South, Taplow, Maidenhead, Berkshire SL6 0PH (GB).		Published With international search report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): CLIFFE, Ian, Anthony [GB/GB]; Wyeth Laboratories, Huntercombe Lane South, Taplow, Maidenhead, Berkshire SL6 0PH (GB). BRIGHTWELL, Christopher, Ian [GB/GB]; Wyeth Laboratories, Huntercombe Lane South, Taplow, Maidenhead, Berkshire SL6 0PH (GB). MANSELL, Howard, Langham [GB/GB]; Wyeth Laboratories, Huntercombe Lane South, Taplow, Maidenhead, Berkshire SL6 0PH (GB). WHITE, Alan, Chapman [GB/GB]; Wyeth Laboratories, Huntercombe Lane South, Taplow, Maidenhead, Berkshire SL6 0PH (GB).			

(54) Title: PIPERAZINE DERIVATIVES

(57) Abstract

Piperazine derivatives of formula (I) and their pharmaceutically acceptable salts are 5-HT_{1A} binding agents and may be used, for example, as anxiolytics. In the formula, A is an optionally substituted alkylene chain, Z is a specified bicyclic nitrogen containing heteroaromatic radical, R is hydrogen or lower alkyl, R¹ is aryl or aryl(lower)alkyl and R² and R³ are hydrogen or specified organic radicals or -NR²R³ represents a heterocyclic ring.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

PIPERAZINE DERIVATIVES


5

This invention relates to piperazine derivatives, to processes for their preparation, to their use and to pharmaceutical compositions containing them. The novel compounds act on the central nervous system by binding to 5-HT receptors (as more fully explained below) and hence can be used as medicaments for treating humans and other mammals.

10

GB 2230781-A and GB 2230780-A disclose some piperazine derivatives which exhibit 5-HT_{1A} receptor affinity.

15 The novel compounds of the invention are those of the general formula



(I)

20 and the pharmaceutically acceptable acid addition salts thereof.

In formula (I)

A is an alkylene chain of 1 or 2 carbon atoms optionally substituted by one or more lower alkyl groups,

25

Z is a bicyclic nitrogen containing heteroaromatic radical selected from the group consisting of optionally substituted indolyl, isoindolyl, quinoliny, isoquinoliny, indazolyl and benzotriazolyl,

30

R represents hydrogen or one or two same or different lower alkyl groups,

R¹ is an aryl radical or an aryl(lower)alkyl radical,

-2-

R² is hydrogen or lower alkyl and

R³ is hydrogen, an alkyl group of 1 to 10 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, cycloalkyl(lower)alkyl, aryl or aryl(lower)alkyl

5

or R² and R³ together with the nitrogen atom to which they are both attached represent a saturated heterocyclic ring which may contain a further hetero atom (eg an azetidino, pyrrolidino, piperidino, hexahydroazepino, heptamethyleneimino, morpholino or piperazino ring which may be optionally substituted by lower alkyl, aryl or aryl(lower)alkyl).

10

The term "lower" as used herein means that the radical referred to contains 1 to 6 carbon atoms. Preferably such radicals contain 1 to 4 carbon atoms. Examples of alkyl radicals are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl and isopentyl. When R³ is an alkyl radical a particularly preferred radical is a tertiary alkyl radical such as tert-butyl. Examples of cycloalkyl groups of 3 to 12 carbon atoms include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. Cycloalkyl groups also include bicyclic, tricyclic and tetracyclic groups eg adamantyl.

20

When used herein "aryl" means an aromatic radical having 6 to 12 carbon atoms (e.g. phenyl or naphthyl) which optionally may be substituted by one or more substituents commonly used in medicinal chemistry, eg substituents such as lower alkyl, lower alkoxy, lower alkylthio, halogen, trifluoromethyl, nitro, carbalkoxy, carboxamido, cyano, amino, (lower)alkylamino and di(lower)alkylamino.

25

Examples of aryl(lower)alkyl include, for example, benzyl in which the phenyl group may be substituted as defined above.

The bicyclic nitrogen containing heteroaromatic radical Z may be unsubstituted or substituted by, for example, one or more same or different substituents selected from lower alkyl, halogen, halo(lower)alkyl (eg trifluoromethyl), nitro, nitrile, oxo, hydroxy, (lower)alkoxy, hydroxy(lower)alkyl, (lower)-alkoxy(lower)alkyl, lower alkanoyloxy(lower)alkyl, lower alkylcarbonyl, (lower)alkylcarbonyl(lower)alkyl, carboxamido, (lower)alkoxy carbonyl, amino, (lower)alkylamino or di(lower)alkylamino. A particularly preferred substituent is (lower)alkoxycarbonyl e.g. -CO.O.CH₃.

35


-3-

Where Z is an optionally substituted indolyl or indazolyl radical, the radical is preferably linked via position 4 or 7 (most preferably via position 4) to the nitrogen atom of the piperazine ring. When Z is an optionally substituted quinoliny or isoquinoliny radical, the radical is preferably linked via position 5 or 8 (most preferably via position 5) to the nitrogen atom of the piperazine ring.

Examples of preferred compounds are

- 10 (A) those in which A is ethylene
- (B) those in which R² is hydrogen and R³ is tert-alkyl or cycloalkyl
- (C) those in which NR²R³ represents a piperidino or hexahydroazepino ring
- 15 (D) those in which Z is optionally substituted indolyl, particularly indol-4-yl optionally substituted by, for example methoxycarbonyl in, for example the 2-position
- 20 (E) those in which R is hydrogen
- (F) those in which R¹ is phenyl


The compounds of the invention may be prepared by methods known in the art from known starting or starting materials that may be prepared by conventional methods. One method comprises alkylation of a piperazine compound of formula



(II)

30

(where Z and R are as defined above) with an alkylating agent providing the group



(III)

-5-

of a coupling agent such as 1,1'-carbonyldiimidazole, iso-butylchloroformate or diphenylphosphinyl chloride.

5 The acids of formula (VII) may be prepared by methods known in the art eg from the piperazine derivatives of formula (II). For example a piperazine derivative of formula (II) may be reacted with an acid of formula



10 by means of a Michael Reaction.

The processes described above may be carried out to give a compound of the invention in the form of a free base or as an acid addition salt. If the compound of the invention is obtained as an acid addition salt, the free base can be obtained by basifying a
15 solution of the acid addition salt. Conversely, if the product of the process is a free base an acid addition salt, particularly a pharmaceutically acceptable acid addition salt, may be obtained by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds.

20

Examples of acid addition salts are those formed from inorganic and organic acids, such as sulphuric, hydrochloric, hydrobromic, phosphoric, tartaric, fumaric, maleic, citric, acetic, formic, methanesulphonic, p-toluenesulphonic, oxalic and succinic acids.

25 The compounds of the invention contain one or more asymmetric carbon atoms, so that the compounds can exist in different stereoisomeric forms. All stereoisomeric forms are included within the invention. The compounds can be, for example, racemates or optically active forms. The optically active forms can be obtained by resolution of the racemates or by asymmetric synthesis.

30

The compounds of the present invention possess pharmacological activity. In particular, they act on the central nervous system by binding to 5-HT receptors, particularly receptors of the 5-HT_{1A} type. In general, the compounds selectively bind to receptors of the 5-HT_{1A} type to a much greater extent than they bind to other
35 receptors such as α_1 . The compounds can be used for the treatment of CNS disorders, such as anxiety in mammals, particularly humans. They may also be useful as

-6-

antidepressants, hypotensives and as agents for regulating the sleep/wake cycle, feeding behaviour and/or sexual function and for treating cognition disorders.

5 The compounds of the invention are tested for 5-HT_{1A} receptor binding activity in rat hippocampal membrane homogenate by the method of B S Alexander and M D Wood, J Pharm Pharmacol, 1988, 40, 888-891. The compound of Example 5, which is a representative compound of the invention, had an IC₅₀ of 0.72 nM.

10 The compounds are also tested for 5-HT_{1A} receptor antagonism activity in a test involving the antagonism of 8-hydroxy-2-(di-n-propylamino)-tetralin (8-OH DPAT) syndrome in the rat. The compound of Example 5 exhibited a median effective dose of 0.3 mg/kg s.c.

15 The invention also provides a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable acid addition salt thereof in association with a pharmaceutically acceptable carrier. Any suitable carrier known in the art can be used to prepare the pharmaceutical composition. In such a composition, the carrier is generally a solid or liquid or a mixture of a solid or liquid.

20 Solid form compositions include powders, granules, tablets, capsules (eg hard and soft gelatine capsules), suppositories and pessaries. A solid carrier can be, for example, one or more substances which may also act as flavouring agents, lubricants, solubilisers, suspending agents, fillers, glidants, compression aides, binders or tablet-disintegrating agents; it can also be an encapsulating material. In powders the carrier is a finely
25 divided solid which is in admixture with the finely divided active ingredient. In tablets the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99%, eg from 0.03 to 99%, preferably 1 to 80% of the active ingredient. Suitable solid carriers include, for example, calcium
30 phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

35 The term "composition" is intended to include the formulation of an active ingredient with encapsulating material as carrier to give a capsule in which the active ingredient (with or without other carriers) is surrounded by the carrier, which is thus in association with it. Similarly cachets are included.

Liquid form compositions include, for example, solutions, suspensions, emulsions, syrups, elixirs and pressurised compositions. The active ingredient, for example, can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fats. The liquid carrier can contain other suitable pharmaceutical additives such as solubilisers, emulsifiers, buffers, preservatives, sweeteners, flavouring agents, suspending agents, thickening agents, colours, viscosity regulators, stabilisers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (particularly containing additives as above, eg cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols, eg glycerol and glycols) and their derivatives, and oils (eg fractionated coconut oil and arachis oil). For parenteral administration the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used in sterile liquid form compositions for parenteral administration.

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. When the compound is orally active it can be administered orally either in liquid or solid composition form.

Preferably the pharmaceutical composition is in unit dosage form, eg as tablets or capsules. In such form, the composition is sub-divided in unit dose containing appropriate quantities of the active ingredient; the unit dosage forms can be packaged composition, for example packeted powders, vials, ampoules, prefilled syringes or sachets containing liquid. The unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form. The quantity of the active ingredient in unit dose of composition may be varied or adjusted from 0.5 mg or less to 750 mg or more, according to the particular need and the activity of the active ingredient.

The following Examples illustrate the invention

-8-

Example 1Methyl pyruvate-3-nitrophenylhydrazone

5 A solution of methylpyruvate (31.8 g, 0.312 mol) in water-ethanol (10:1, 100 ml) was slowly added to a hot solution of 3-nitrophenylhydrazine hydrochloride (59.2 g, 0.312 mol) and sodium acetate (28.0 g, 0.34 mol) in water and ethanol (5:7, 1200 ml). The mixture was stirred at room temperature for 2 h during which time a heavy yellow
10 precipitate developed. The precipitate was filtered, washed with cold water (2000 ml), and dried to give the product (66 g) as a yellow solid.

Example 2Methyl 4-nitroindole-2-carboxylate

The product of Example 1 (66 g, 0.278 mol) and polyphosphoric acid (1000 g) were stirred slowly as the temperature was raised to 90° C, at which point an exotherm occurred. The reaction mixture turned orange/brown and its viscosity increased .
20 Further polyphosphoric acid (200 g) was added and the reaction mixture was stirred vigorously keeping the temperature below 100° C by ice cooling. After 1 h, the mixture was poured into an ice/water mixture (5000 ml) and extracted into ethyl acetate (2 x 1500 ml). The extracts were washed with dilute aqueous sodium hydroxide (1000 ml), dried (MgSO₄), and evaporated in vacuo to give a mixture of methyl 4-nitroindole-
25 2-carboxylate and methyl 5-nitroindole-2-carboxylate (total weight=45.6 g) as a grey yellow solid. This isomeric mixture was used without separation as the starting material in Example 3.

Example 3

30

Methyl 4-aminoindole-2-carboxylate

Ammonium formate (63 g, 1.00 mol) was added cautiously to a suspension of the product of Example 2 (45.6 g, 0.207 mol) and 10% palladium on charcoal (5 g) in
35 methanol (700 ml) under an argon blanket at 45° C. The mixture was stirred for 4 h, cooled to room temperature, and after 18 h filtered and evaporated in vacuo to give a dark brown semi-solid. This was taken up into acetonitrile (500 ml), filtered (to remove

-9-

ammonium formate) and evaporated in vacuo to give a dark brown gum. Purification by column chromatography [silica; ethyl acetate-hexane (1:10 to 1:1)] gave the product (9.5 g) as a grey solid.

5

Example 4

Methyl 4-(1-piperazinyl)indole-2-carboxylate

10 A mixture of the product of Example 3 (9.5 g, 0.051 mol), bis(2-chloroethyl)amine hydrochloride (10.6 g, 0.06 mol), and potassium carbonate (8.3 g, 0.06 mol) in anhydrous dimethylformamide (30 ml) was heated at 110° C under argon. Further addition of bis(2-chloroethyl)amine hydrochloride (5.5 g, 0.031 mol) and potassium carbonate (4.2 g, 0.031 mol) were made after 7 h and 16 h. After another 6 h, the
15 reaction mixture was poured into dilute potassium carbonate solution (400 ml) and extracted with ether (3 x 300 ml) and ethyl acetate (2 x 250 ml). The extracts were combined, dried (MgSO₄), and washed with water (2 x 200 ml). Evaporation of the solvent in vacuo gave a solid which was recrystallised from ether to give the product (5.0 g) as off-white crystals, m.p. 177-180° C.

20

Example 5

4-[4-(4-Azepan-1-yl)-4-oxo-3-phenylbutyl]piperazin-1-yl]-1H-indole-2-carboxylic acid methyl ester

25

The product of Example 4 (1.30 g, 0.005 mol), 1-azepan-1-yl-4-chloro-2-phenylbutan-1-one (1.50 g, 0.005 mol), triethylamine (0.79 g, 0.0079 mol), and potassium iodide (0.2 g) were suspended in anhydrous dimethyl formamide (30 ml), stirred under argon for 3 h at 100° C, and poured into dilute potassium carbonate (300 ml). The mixture
30 was extracted into ether (3 x 100 ml) and treated with excess dilute aqueous hydrogen chloride. The precipitate was filtered, washed with ether (3 x 100 ml), taken up into dilute sodium hydroxide solution (100 ml), and extracted into ether (3 x 100 ml). The extracts were dried (MgSO₄), and evaporated in vacuo to give a yellow crystalline glass. Purification by chromatography [alumina; ethyl acetate - hexane (2:3)] gave the
35 product free base. The dihydrochloride salt of the product (1.35 g) was precipitated as a colourless solid by the addition of ethereal hydrogen chloride to a solution of the free base in ethyl acetate, m.p. 152-168° C (Found: C, 57.4; H, 7.45; N, 8.7.

-10-

$C_{30}H_{38}N_4O_3 \cdot 2HCl \cdot 3H_2O$ requires C, 57.2; H, 7.4; N, 8.9%).

Example 6

5 1-Azepan-1-yl-2-phenyl-4-(4-quinolin-5-yl-piperazin-1-yl)-butan-1-one

A mixture of 5-(1-piperazinyl)quinoline (932 mg, 4.37 mmol), potassium iodide (725 mg, 4.37 mmol), N,N-diisopropylamine (0.9 ml, 5.2 mmol), and 1-azepine-1-yl-4-chloro-2-phenylbutan-1-one (1.22 g, 4.37 mmol) in dimethylformamide (20 ml) was
10 heated at 110° C for 16 h, cooled to room temperature, treated with water (150 ml), and extracted with ethyl acetate (2 x 50 ml). The extracts were washed with water (50 ml), dried (Na_2SO_4), and evaporated in vacuo. The residue was purified by chromatography [alumina; ethyl acetate-hexane (2:3)] to give the free base of the product (1.10 g). The dihydrochloride salt was a bright yellow crystalline solid, m.p. 216-219° C (Found: C,
15 63.4; H, 7.65; N, 10.15. $C_{29}H_{36}N_4O \cdot 2HCl \cdot 1\frac{1}{4}H_2O$ requires C, 63.1; H, 7.4; N, 10.15%).

Example 7

20 1-Azepan-1-yl-2-phenyl-4-(4-quinolin-8-yl-piperazin-1-yl)-butan-1-one

This compound was prepared from 8-(1-piperazinyl)-quinoline using the method described in Example 6. The one and a half hydrochloride salt of the product was obtained as sandy crystals, m.p. 223-226° C (Found: C, 64.7; H, 7.7; N, 10.2.
25 $C_{29}H_{36}N_4O \cdot 1\frac{1}{2}HCl \cdot 1\frac{1}{2}H_2O$ requires C, 64.7; H, 7.6; N, 10.4%).

Example 8

30 4-[4-(4-Azepan-1-yl-4-oxo-3R or 3S-phenylbutyl)-piperazin-1-yl]-1H-indole-2-carboxylic acid methyl ester

Racemic 4-[4-(4-azepan-1-yl-4-oxo-3-phenylbutyl)piperazin-1-yl]-1H-indole-2-carboxylic acid methylester was resolved into the R and S enantiomers by high performance liquid chromatography on a chiral cell OD-R column (4.6 x 250 mm)
35 using 60:40 mixture of acetonitrile : 0.5M sodium perchlorate buffer (pH 5.8). Detection was by U.V. (λ 254 nm) flow rate 1ml/minute. Under the conditions used the R enantiomer had a retention time of 32 min, and the S 4/min.

-11-

After work up the S-enantiomer was isolated as its dihydrochloride hemihydrate. mp 148-149°C. Analysis Found: C, 61.8; H, 7.4; N, 9.5%. $C_{30}H_{38}N_4O_3 \cdot 2HCl \cdot 0.5H_2O$ requires C, 61.6; H, 7.1; N, 9.6%.

5

The R enantiomer was isolated as its dihydrochloride three-quarter hydrate mp. 148-149°C. Found: C, 61.5; H, 7.2; N, 9.5%. $C_{30}H_{38}N_4O_3 \cdot 2HCl \cdot 0.75H_2O$ requires C, 61.2; H, 7.1; N, 9.5%.

10

Example 9

1-Azepan-1-yl-4-[4-(1H-indol-4-yl)piperazin-1-yl]-2-phenylbutan-1-one

- 15 A suspension of 4-aminoindole (2.30 g, 0.017 moles), bischloroethylamine hydrochloride (3.49 g, 0.02 moles) and potassium carbonate (5.41 g, 0.02 moles) in butanol (50 ml) was heated at reflux under argon for 7h. Further bischloroethylamine hydrochloride (0.70 g, 0.04 moles) and potassium carbonate (1.1 g 0.04 moles) was added, then the suspension was heated at reflux for 18 h.
- 20 The solvent was evaporated in vacuo to give a brown gummy residue which was taken up into dilute hydrochloric acid (200 ml), washed with ethyl acetate (2 x 100 ml), made basic with potassium carbonate, extracted into dichloromethane (3 x 50 ml), dried ($MgSO_4$) then evaporated in vacuo to give a white solid which was triturated with ethyl acetate and dried to give 4-(1-piperazinyl)indole (0.78 g,
- 25 0.0038 moles) as white crystals.

- A solution of the above product (0.70 g, 0.0035 moles), 1-azepine-1-yl-4-chloro-2-phenylbutan-1-one (1.12 g, 0.0040 moles), triethylamine (0.49 ml, 0.0035 moles) plus potassium iodide (0.2 g) in dimethylformamide (20 ml) was heated at reflux
- 30 for 2 h then stirred at room temperature for 18 h under argon. The solvent was evaporated in vacuo to give a yellow oil which was taken up into ethyl acetate (50 ml), washed with dilute sodium carbonate solution (2 x 50 ml), dried ($MgSO_4$) then evaporated in vacuo to give a yellow oil. This was purified on an alumina column, eluting firstly with 30% ethyl acetate in hexane then with ethyl acetate
- 35 alone to give the title compound (1.0 g) as a clear oil.

-12-

The above product was taken up into ethyl acetate then precipitated with ethereal hydrogen chloride to give the hydrochloride (0.75 g, 0.0014 moles) as a white solid m.p. 174-177°C. (Found: C, 62.4; H, 7.6 N, 10.3%. $C_{28}H_{36}N_4O \cdot 2HCl \cdot 1.25H_2O$ requires C, 62.3; H, 7.6; N, 10.4%).

5

Example 10

1-Azepan-1-yl-4-[4-(1H-indol-7-yl)piperazinyl-1-yl]-2-phenylbutan-1-one

- 10 A suspension of 7-aminoindole (4.10 g, 0.031 moles), bischloroethylamine hydrochloride (6.23 g, 0.035 moles) and potassium carbonate (9.66 g, 0.035 moles) in butanol (100 ml) was heated at reflux under argon for 6h. Further bischloroethylamine hydrochloride (3.0 g, 0.016 moles) and potassium carbonate (4.5 g, 0.016 moles) was added, then the suspension was heated at reflux under
15 argon for 18 h. The solvent was evaporated in vacuo to give a deep purple residue which was taken up into dilute hydrochloric acid (300 ml), washed with ethyl acetate (2 x 100 ml), made basic with sodium hydroxide, extracted into dichloromethane, filtered, dried ($MgSO_4$) then evaporated in vacuo to a yellow oil. This was crystallised twice from ether to give 7-(1-piperazinyl)indole (2.07 g,
20 0.010 moles) as a grey/white solid.

- A solution of 7-(1-piperazinyl)indole (1.00 g, 0.005 moles), 1-azepine-1-yl-4-chloro-2-phenylbutan-1-one (1.50 g, 0.0055 moles), triethylamine (0.50 g, 0.005 moles) plus potassium iodide (0.2 g) in dimethylformamide (30 ml) was heated at
25 reflux under argon for 6 h. The solvent was evaporated in vacuo to give a brown residue which was taken up into dilute sodium carbonate solution (100 ml), extracted into ethyl acetate (2 x 100 ml), dried $MgSO_4$ then evaporated in vacuo to give a brown solid which was triturated with ethyl acetate (2 x 50 ml) and isopropanol (3 x 50 ml) to give the title compound (0.9 g) as a white solid.

30


The above product was taken up into dichloromethane (50 ml), then precipitated with ethereal hydrogen chloride to give the hydrochloride as a white solid, m.p. 138-145°C. (Found: C, 65.7; h, 7.8; N, 10.8% . $C_{28}H_{36}N_4O \cdot 1.75HCl \cdot 0.25H_2O$ requires: C, 65.6; H, 7.52; N, 10.9%.)

35

-13-

CLAIMS

1. A compound of formula



or a pharmaceutically acceptable acid addition salt thereof.

wherein

A is an alkylene chain of 1 or 2 carbon atoms optionally substituted by one or more lower alkyl groups,

Z is a bicyclic nitrogen containing heteroaromatic radical selected from the group consisting of optionally substituted indolyl, isoindolyl, quinoliny, isoquinoliny, indazolyl and benzotriazolyl,

R represents hydrogen or one or two same or different lower alkyl groups,

R¹ is an aryl radical or an aryl(lower)alkyl radical,

R² is hydrogen or lower alkyl and


R³ is hydrogen, an alkyl group of 1 to 10 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, cycloalkyl(lower)alkyl, aryl or aryl(lower)alkyl

or R² and R³ together with the nitrogen atom to which they are both attached represent a saturated heterocyclic ring which may contain a further hetero atom.

2. A compound as claimed in claim 1 in which A is ethylene.


-14-

3. A compound as claimed in claim 1 or 2 in which R^2 is hydrogen and R^3 is tert-alkyl or cycloalkyl.
4. A compound as claimed in claim 1 or 2 in which NR^2R^3 represents a piperidino or hexahydroazepino ring.
5. A compound as claimed in any one of claims 1 to 4 in which Z is optionally substituted indolyl.
6. A compound as claimed in claim 1 which is 4-[4-(4-azepan-1-yl-4-oxo-3-phenylbutyl)piperazin-1-yl]-1H-indole-2-carboxylic acid methyl ester, 1-azepan-1-yl-2-phenyl-4-(4-quinolin-5-yl-piperazin-1-yl)-butan-1-one, 1-azepan-1-yl-2-phenyl-4-(4-quinolin-8-yl-piperazin-1-yl)-butan-1-one, 4-[4-(4-azepan-1-yl-4-oxo-3-phenylbutyl)piperazin-1-yl]-1H-indole-2-carboxylic acid methyl ester, 1-azepan-1-yl-4-[4-(1H-indol-4-yl)piperazin-1-yl]-2-phenylbutan-1-one and 1-azepan-1-yl-4-[4-(1H-indol-7-yl)piperazinyl-1-yl]-2-phenylbutan-1-one or a pharmaceutically acceptable acid addition salt thereof.
7. A process for preparing a compound claimed in claim 1 which comprises
 - (a) alkylating a piperazine compound of formula



(II)

(where Z and R are as defined in claim 1) with an alkylating agent providing the group



(III)

(where A, R^1 , R^2 and R^3 have the meanings given in claim 1)


or

- (b) acylating an amine of formula

-15-



(where R^2 and R^3 are as defined in claim 1) with an acid of formula



(where Z, A, R and R^1 are as defined in claim 1) or with an acylating derivative thereof

or

(c) resolving a compound claimed in claim 1 into an enantiomer

or

(d) converting a base claimed in claim 1 into a pharmaceutically acceptable salt or converting a pharmaceutically acceptable salt into the free base.

8. A pharmaceutical composition comprising a compound claimed in claim 1 in association with a pharmaceutically acceptable carrier.

9. A compound claimed in claim 1 for use as a pharmaceutical.

10. A compound claimed in claim 1 for use as a 5-HT_{1A}-antagonist.

INTERNATIONAL SEARCH REPORT

 Interna' 1 Application No
 PCT/GB 93/02660

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 5 C07D209/42 C07D215/38 C07D215/40 C07D209/08 A61K31/40
 A61K31/47 A61K31/55

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 395 312 (JOHN WYETH & BROTHER LIMITED) 31 October 1990 *see especially claims 3,4 and 5, examples and description, page 3, lines 15-23* & GB,A,2 230 781 (JOHN WYETH & BROTHER LTD) cited in the application ---	1-10
Y	EP,A,0 395 313 (AMERICAN HOME PRODUCTS CORP AND JOHN WYETH & BROTHER LTD.) 31 October 1990 *see examples and whole document* & GB,A,2 230 780 (JOHN WYETH & BROTHER LTD) cited in the application ---	1-10

-/--

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

10 March 1994

Date of mailing of the international search report

25. 03 94

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Scruton-Evans, I

INTERNATIONAL SEARCH REPORT

Internat 1 Application No

PCT/GB 93/02660

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,92 03426 (NOVO NORDISK A/S) 5 March 1992 *see definition of R1 and examples 30-33,37,38,40* ---	1-10
Y	EP,A,0 512 755 (JOHN WYETH & BROTHER LTD.) 11 November 1992 *see definition of R1* ---	1-10
P,Y	WO,A,93 11122 (JOHN WYETH & BROTHER LTD.) 10 June 1993 *see definition of Z* ---	1-10
A	MEDICINAL CHEMISTRY RESEARCH vol. 2, no. 2 , 1992 , BOSTON pages 88 - 95 M. EL-BERMAWY ET AL '4-[4-(1-noradamantane carboxamido)butyl]-1-(2-methoxyphenyl)pipe razine: A high affinity 5-HT1A selective agent' -----	1-10

INTERNATIONAL SEARCH REPORT

information on patent family members

Intern: Application No

PCT/GB 93/02660

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0395312	31-10-90	AU-B- 619677	30-01-92
		AU-A- 5377890	25-10-90
		AU-B- 619678	30-01-92
		AU-A- 5377990	25-10-90
		EP-A- 0395313	31-10-90
		GB-A, B 2230780	31-10-90
		GB-A, B 2230781	31-10-90
		GB-A, B 2255976	25-11-92
		JP-A- 3011059	18-01-91
		JP-A- 3020263	29-01-91
		US-A- 4921958	01-05-90
		US-A- 4988814	29-01-91
GB-A-2230781	31-10-90	AU-B- 619677	30-01-92
		AU-A- 5377890	25-10-90
		AU-B- 619678	30-01-92
		AU-A- 5377990	25-10-90
		EP-A- 0395312	31-10-90
		EP-A- 0395313	31-10-90
		GB-A, B 2230780	31-10-90
		GB-A, B 2255976	25-11-92
		JP-A- 3011059	18-01-91
		JP-A- 3020263	29-01-91
		US-A- 4921958	01-05-90
		US-A- 4988814	29-01-91
EP-A-0395313	31-10-90	AU-B- 619677	30-01-92
		AU-A- 5377890	25-10-90
		AU-B- 619678	30-01-92
		AU-A- 5377990	25-10-90
		EP-A- 0395312	31-10-90
		GB-A, B 2230780	31-10-90
		GB-A, B 2230781	31-10-90
		GB-A, B 2255976	25-11-92
		JP-A- 3011059	18-01-91
		JP-A- 3020263	29-01-91
		US-A- 4921958	01-05-90
		US-A- 4988814	29-01-91
		CA-A- 2015033	22-10-90

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 93/02660

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-2230780	31-10-90	AU-B- 619677	30-01-92
		AU-A- 5377890	25-10-90
		AU-B- 619678	30-01-92
		AU-A- 5377990	25-10-90
		EP-A- 0395312	31-10-90
		EP-A- 0395313	31-10-90
		GB-A, B 2230781	31-10-90
		GB-A, B 2255976	25-11-92
		JP-A- 3011059	18-01-91
		JP-A- 3020263	29-01-91
		US-A- 4921958	01-05-90
		US-A- 4988814	29-01-91
		CA-A- 2015033	22-10-90
WO-A-9203426	05-03-92	AU-A- 8441391	17-03-92
		CA-A- 2089768	25-02-92
		EP-A- 0544765	09-06-93
		JP-T- 6500546	20-01-94
		US-A- 5246935	21-09-93
EP-A-0512755	11-11-92	AU-B- 645681	20-01-94
		AU-A- 1524192	05-11-92
		GB-A- 2255337	04-11-92
		JP-A- 5170743	09-07-93
WO-A-9311122	10-06-93	AU-A- 2953792	28-06-93
		GB-A- 2262093	09-06-93